

Libyan International Medical University



Faculty of dentistry

Comprehensive Case Presentation

Presented by:

Nora Ali Alabdly

Personal Data:-

Patient's Name: N.A.E.

D.O.B: 1/1/1987.

Gender: female.

Occupation: House wife

سيدي حسين : Address

Nationality: Libyan.

Chief complaint:-

Pain in the upper right teeth since 2 day.

History of chief complaint:-

Sever localized spontaneous throbbing pain, started 2 months ago as a mild intermittent pain and became sever 2day, it increases on biting, lasts for hours sometimes it subsides by itself and in other times gets less with analgesics (Ibuprofen 200 mg 2 tab /d), disturbs with sleep, and it's associated with headache.

Past & current medical history:-

The patient is medically fit.

Allergic History: -

No history of allergies.

Past dental history:-

Last dental visit from 2 years ago Extractions without complication.

Practice:

Brushing: irregular with soft tooth brush.

Brushing method: Horizontal.

Social and Family History:

- single, nonsmoker, non alcohol consumer, consumes frequent sugary food & sweet drink drinks water frequently (2L/day).
- Diabetes {Mother's side}.

Clinical Examinations :-

Extra Oral Examinations:-

1-Face symmetry: symmetrical. 2.Skin: Normal color, no scars. 3.TMJ : Average mouth opening, No tenderness, clicking, deviation 4.Lymph nodes: Not palpable.

5.Lips: competent

Intra Oral Examinations:

Soft tissues Examinations:-

Oral mucosa -palate- Floor of the mouth and tongue are

normal













Intra Oral Examinations:

Hard tissues Examinations:-

Rt Lateral View



Lf Lateral View



Front View



Hard tissues Examinations:-

Maxillary Arch:-



8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8
M	M	D	D	D							D			D	M

Hard tissues Examinations:-

Mandibular Arch:-





D	D	D	D	D								D	D	M	D
8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8

Gingival status:-

	Max . Right posterior	Max . Anterior	Max . Left Posterior
Color	Reddish-pink	Reddish-pink	Reddish-pink
Contour	Rounded margin + blunt IDP.	Rounded margin + blunt IDP.	Rounded margin + blunt IDP.
Size	Slightly enlarged.	Slightly enlarged.	Slightly enlarged.
Consistency	Edematous.	Edematous.	Edematous.
Stippling	+ve	+ve	+ve
Position	Coronal to CEJ.	Coronal to CEJ.	Coronal to CEJ.
Bleeding	+	+	+
Exudation	-	-	-

Gingival status:-

	Man . Right posterior	Man . Anterior	Man . Left Posterior
Color	Reddish-pink	Reddish-pink	Reddish-pink
Contour	Rounded margin + blunt IDP	Rounded margin + blunt IDP.	Rounded margin + blunt IDP.
Size	enlargement.	enlargement.	enlargement.
Consistency	Edematous.	Edematous.	Edematous.
Stippling	+	+	+
Position	Apical to CEJ.	Coronal to CEJ.	Coronal to CEJ.
Bleeding	+	+	+
Exudation	-	-	-

Periodontal Status

- General probing depth within normal limit 2-3 mm with exception 6532 5

Between 4-5 mm
Attachment loss 6532
876
Between 2-3 mm
Gingiva recession 876
No furcation involvement
And no mobility of teeth

Adequacy of Attached Gingiva:- Adequate.


Fremitus test: no trauma from occlusion

Frenal Attachment:- Normal.

Oral Hygiene:-

Oral hygiene index-simplified:

16	11	26
2	1	2
2	2	2
46	31	36



	ACCURATE STREET		
y	16	11	26
	1	1	1
	1	1	1
	46	31	36

Plaque index= 11/6= 1.8

OHI-S=1.8+1=2.8

Calculus index= 6/6= 1



Dietary Analysis:-

		THE RESIDENCE OF THE PROPERTY	() () () () () ()	PER SHARE A SUM OF STREET
1	Patient Name: Occupation:	Gend Gend	aress.	
	نوح الوجية	الماكولات والمشروبات	الباعة	الأيام
8	وعور	do orio cuto upo		
	عبداء	alo_ (my abelia agosa		N.
	Syper	Jee - 43-15 gin	6:30	
	- اشد	بيتزاء ـ فاكمه _ ماء	10:00	
Š.	1960	دله - بعد عنال تسق	11:00	
	- laie	طاحين دماج - سلاطه: طارعه ما	2:00	3
	2 pe	الماء الماء الماء		3
		زنجر - سبس - ماد		
9		16. Culsel 4615 - 200 - (1)	11:30	
		6- Come - chee- Cosus		
		ماء		
		سلاطة فواجة		
		سانهوتش _ بيس _ ماد. فاة	10 = 30	
			1973 88	

Gnvesligations

Orthopantamograph (OPG)




Showing maxillary and mandibular teeth with mild bone loss Missing teeth

Missing teeth

Investigations:-

Bitewing





Final Diagnosis

- Chronic periapical periodontitis with acute exacerbation
- 4
- -localized mild to moderate chronic periodontitis.
- Chronic Irreversible pulpitis 5 4
- -Dental caries 6 7 87654 568

Treatment plane :-

Emergency phase:- Extraction

```
4
```

Phase I:-

- OHI :-Patient motivation and education.
- Scaling & root planning.

Re-evaluation of Phase I

Phase II therapy:-

Extraction of non-restorable teeth



Phase III therapy:-

Phase IV:-

Follow up & maintenance.

Informed Consent:-

الجامعة الليبية الدولية للعلوم الطبيه كلية طب وجراحة الفع والأسنان موافقة للأذن بالعلاج ويكامل ارادتي أنفي أعطى موافقتي اللفضيية والكتابية للبدء والاستمرار في العلاج بعيادات كلية طب الأستان بالح وقدتم شرح التشخيص المرضمي لحالتي والخيارات المختلفة المتوفرة لمعالجتها والأعراض الجانبية والمخاطر التي قد تصاحب العلاج والمشاكل التي قد تنتج عن عدم العلاج والعدة التقريبية اللازمة للانتهاء منه والتكلفة الاجماليه للعلاج وققا للرسوم الرمزية المقررة كما اتفهم ان هذا العلاج قد تم تنفيذه من قبل طالب أو طبيب امتياز تحت الاشراف المباشر لأعضاء هيئة التدريس بالكلية

Emergency phase:-

Extraction of non-restorable teeth







PHASE 1 THERAPY:-

- -OHI:-Patient motivation and education.
- Scaling and root planning.
- Diet counseling.

Fluoride application





Fluoride application:-


Reevaluation of PHASE I:-

- -Patient showed significant improvement of oral hygiene.
 - -Improved gingival status.
 - -Reduction of pockets depth.

Reevaluation of PHASE I:-

Oral hygiene index-simplified:

11	26
0	1
0	0
31	36
	0



7	16	11	26
	0	0	0
	0	0	0
A CONTRACTOR OF THE PARTY OF TH	46	31	36
N. Constitution of the Con	46	31	36

Plaque index=2/6 = 0.33

Calculus index= 0/6 = 0

$$OHI-S=0.33+0=0.33$$



PHASE II: -

Extraction of non-restorable teeth

8






PHASE III: -

RCT 5 4






PHASE III : -

$$RCT = \frac{5}{}$$







PHASE III: - Restoration 6 indirect pulp capping, Ca(ho)2, 6 GIC, composite restoration shade 2









PHASE III: -

Restoration







PHASE III:-

Restoration










PHASE III: -

Restoration





PHASE III:-

Restoration








56





PHASE III:-

Restoration







PHASE III:-

Restoration







PHASE III: -

Restoration of endodontic treated teeth by using ready made fiber Post and composite

Core 5 | 4





Post placement and composite core build up









Preparation of tooth—4—Final impression



Temporary crown fabrication





Composite Try-in on the cast







Composite Try-in

Checking the margins



Checking the occlusion with articulating paper



Shade Selection





E- Max crown Try-in on the cast



E- Max crown Try-in



Checking the occlusion with articulating paper

Checking the margins









Final cementation of E- Max crown by self adhesive resin cement







PHASE III: -

Restoration of endodontic treated teeth by using ready made fiber Post and composite Core 5





Post placement and composite core build up





Preparation of tooth



Final impression




Temporary crown fabrication





Composite Try-in on the cast





Composite Try-in

Checking the occlusion with articulating paper

Checking the margins





E- Max crown Try-in on the cast










Final cementation of E- Max crown by self adhesive resin cement





Recall visit





Primary impression for upper and lower arch to fabricate RPD



Diagnostic cast with RPD













PHASE IV: -

Follow up and Maintenance

- Recall after 3 months:
- -To evaluate oral hygiene & periodontal health.
- -To evaluate all the restorations & the prosthesis.
- Recall for 6 months for regular dental check ups.



Before and After







Before and After











